

Dividing the cerebral cortex surface Sulci – folds

• Gyri – bulges

Superior temporal gyrus

Functional divisions

- Frontal
 - Future planning, personality, <u>judgment</u>, social <u>behavior</u>
 - Motor cortex
- Temporal
 - Auditory cortex
 - High-level vision
- Parietal
 - Spatial vision
 - Primary sensory cortex
 - Visual-auditory-spatial sensory integration
- Occipital
 - Primary visual cortex


7

12

Limbic system – memory/emotion

Medial area of cerebral cortex

- Hippocampus: memory
- Amygdala: emotion
- Cingulate and parahippocampal gyri





The brain beyond the neocortex

- Brain Stem
 - <u>Conduit</u> for spinal cord and cranial nerves
 - Respiratory and cardiac activity
- Cerebellum
 - Plan, coordinate, modify motor activities



Creative Commons, some right reserved http://en.wikipedia.org/wiki/File:1806_The_Hypothalamus-Pituitary_Complex.jpg





Cortical processing networks

Perception "hierarchy"


- Primary sensory areas capture basic sensory properties, or "features"
- More complex representations in higher sensory areas
- Example: Vision
 - Primary visual cortex in occipital pole
 - Anterior flow of information in ventral ("what") and dorsal ("where") pathways

Creative Commons, some rights reserved http://en.wikipedia.org/wiki/File:Ventral-dorsal streams.svg


Cortical processing networks


Action


- Motor planning and performance is achieved in stages within the frontal lobe
- Motor correction is supervised by the cerebellum




http://en.wikipedia.org/wiki/File:Ventral-dorsal_streams.svg











Complexity of cortical networks

- Lateral connections: collaboration among cortical regions at similar stage of processing
- Multi-level connections: shortcuts between "early" and "late" processing stages





Cortical modules

Groups/"columns" of neurons encoding same property

Subdivisions within cortical region


- Retinotopy
- Body part map
 Homunculus
- Tonotopy

http://www.ncbi.nlm.nih.gov/books/NBK10946/

Recording activity in the brain


- Electrode recordings
 - Invasive
 - Single neurons close together
- Neuroimaging
 - Non-invasive
 - Hundreds-millions of neurons broadly distributed
- Behavior
 - Non-invasive
 - Overall cognition

27




Electroencephalography (EEG) Records electric fields at skull surface ~100 sensors across the head low spatial resolution high (ms) time resolution cheap

Magnetoencephalography (MEG) Records magnetic "dipoles" from deep in brain 300 sources found inside the head low spatial resolution ms time resolution expensive




functional Magnetic Resonance Imaging (fMRI)

- Records concentration of blood oxygen
- 10,000+ cortical cubes (2x2x 2 mm³) relatively high spatial resolution
- Low (0.5-4 seconds) time resolution
- expensive








Tractography: Diffusion Tensor Imaging (DTI

Uses MRI

Track flow of water along axon tracts

• Pick source location, see where water flows to



Each color is a different white matter pathway

https://www.mayfieldclinic.com/PE-fMRI_DTI.HTM

34